PENNSTATE HERSHEY Cancer Institute

Intrathecal Chemotherapy

REVISED: June 2012

| Date/time written | To be | gin | | | |
|---|--------------------------------------|------------------|--------------------|--------------------------|---------|
| Patient's: Heightcr Allergies: ■No Diagnosis | □Yes: | | | | |
| 1. Consent Obtained? ☐ Yes: ☐ Preprinted C ☐ No: Plan: 2. Infusion Room General C | Consent | ted Note | te in Chart | | |
| 3. Premedications: Hydratic | Order Set will be initiate on: tics: | | | | |
| 4. Chemotherapy: | | | | | |
| Patient has Ommaya rese | ervoir | | | | |
| ☐ Methotrexate | _ mg in 3ml of sterile, | preservative-fr | ee normal saline | for intrathecal adminis | tration |
| □ Etoposide n | ng in 3 ml of sterile, pr | eservative-free | normal saline fo | or intrathecal administr | ation |
| ☐ Cytarabine | mg in sterile, preserva | ative-free norm | al saline for intr | athecal administration | |
| ☐ Liposomal cytaral | oine mg vial for | intrathecal adı | ninistration (50 | mg vial is standard dose | e) |
| □ Topotecan | mg in 3 ml of sterile, | preservative-fr | ee normal saline | for intrathecal adminis | tration |
| ☐ Trastuzumab administration | mg in 3 ml of sterile | e, preservative- | free normal salir | e for intrathecal | |
| ☐ Rituximab | mg in 3 ml of sterile, | preservative-fr | ee normal saline | for intrathecal adminis | tration |
| | mg sterile, pr | eservative-free | normal saline fo | r intrathecal administra | ation |
| | mg sterile, pr | eservative-free | normal saline fo | r intrathecal administra | ation |
| | Physicia | an/Preparer's S | ignature | | _ |
| | Attendi | ing signature (r | eanired) | | |



MR CHEMO ORDER