PENNSTATE HERSHEY
Cancer Institute

REVISED December 8, 2010

VIP (germ cell tumor).

Date written _____ To begin _____

J Clin Oncol 1998;16:1287-93.

Patient's: Height_____cm Weight_____kg BSA_____m^2
Allergies: ☐No ☐Yes: _____
Diagnosis _____Metastatic Site _____

Protocol:
☐Yes #_____
☐ No
Source of Drug:
☐Routine supply
☐Protocol supply
☐Other

1. Laboratory Studies:
   ☐ None needed prior to first dose of chemo
   ☐ Additional labs needed prior to first dose chemo:
   CBC w/ diff, CMP prior to chemo, Urinalysis (macroscopic only) Qshift
2. Consent Obtained ?
   ☐ Yes☐ Preprinted Consent ☐ See Dictated Note ☐ Note in Chart
   ☐ No Plan: _____
3. Hydration:
   Normal Saline 1000 mL infuse at 500 mL/hr x 2 hours pre-cisplatin on day 1, then via continuous infusion at 100 mL/hr on days 1-5
4. Premedications:
   Ondansetron 24 mg IV q24hr prior to chemotherapy on Days 1-5
   Dexamethasone 10 mg IV q 24hr prior to chemothern Days 1-5
5. **Chemotherapy dose calculation:**

   Etoposide 75 mg/m^2 = _____mg IV in 500 mL NS over 60 minutes q24hr on days 1-5

   Cisplatin 20 mg/m^2 =_____mg IV in 150 mL NS, infuse over 30 minutes q24h on days 1-5 (do not infuse with Mesna- flush tubing before & after)

   Mesna 120 mg/m^2 = _____mg IV in 50 mL D5W infuse over 15 minutes before ifosfamide on day 1 only

   Ifosfamide 1200 mg/m^2 = _____mg IV in 250 mL NS, infuse over 1 hour q24h on days 1-5

   Mesna 1200 mg/m^2 = _____mg IV in 1L NS, infuse via continuous infusion q24hr on days 1-5.
6. Post-discharge care:
   Dexamethasone 4 mg PO BID on Days 6-8
   Ondansetron 8 mg PO BID on Days 6-8
   Prochlorperazine 10 mg PO Q6hr PRN nausea/vomiting
   Pegfilgrastim 6 mg via subcutaneous injection x 1 dose to be given 24-72 hr post-chemotherapy infusion
   Laboratory studies: CBC Weekly, fax results to #717-531-4969.

MD/Preparer's Signature _____

Attending Physician Signature_____

MR CHEMO ORDER