

Revised: March 2011 <b>Date written</b>		To hegin			FOLFOX 6	(Inpatient <sub>)</sub>
Patient's: Height		_			Protocol:  □Yes #  □ No  Source of Drug: □Routine supply	
1. Laboratory Studies:  None neede Additional	d prior to first do labs needed prio	ose of chemo	o e chemo:		□Protocol supply □Other	d plt < 100 k
	reprinted Conse				■ Note in Chart	
3. ACTIVATE ORD	ER SET CHEM	O: FOLFO	X 6 ADULT			
4. Premedications: Ant	D C	examethaso alcium Gluo	16 mg IV/PO one 20 mg IV/P conate 1 Gm an over 30 min pre an 0 mg PO or IV	nd 8 mEq nd post Oxal		e IV
5. Chemotherapy dose	calculation: 🗖 🛭	Oose Reduce	to% of	f full dose.		
Oxaliplatin 10 (Only compati	0 mg/m <sup>2</sup>	raining solutions	ng IV in 500 m . Flush line with D5	nl D5W or W before and	ver 2 hours, Days 1 after completion of oxaliple	and 15 atin. Avoid saline
	0 mg/m <sup>2</sup> = (Infuse with oxaliplati			ml D5W o	over 2 hours, Days 1	and 15
5-Fluorouracil	$400 \text{ mg/m}^2 = \underline{\hspace{1cm}}$	1	mg IV in 50 m	ıl D5W ov	er 15 min, Days 1 a	and 15
]	il 2400 mg/m <sup>2</sup> = Days 1 and 15 v Patient to be seen	ia infusion	pump.		fusion over 46 hour	'S
		MD	/Preparer's Si	gnature _		
		Attendi	ng Physician S	ignature_		

